

Single-Atom Demonstration of the Quantum Landauer Principle

L. L. Yan,¹ T. P. Xiong,^{1,2} K. Rehan,^{1,2} F. Zhou,^{1,*} D. F. Liang,^{1,3} L. Chen,¹ J. Q. Zhang,¹
W. L. Yang,^{1,†} Z. H. Ma,⁴ and M. Feng^{1,3,5,6,‡}

¹*State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China*

²*School of Physics, University of the Chinese Academy of Sciences, Beijing 100049, China*

³*Synergetic Innovation Center for Quantum Effects and Applications (SICQEA), Hunan Normal University, Changsha 410081, China*

⁴*Department of Mathematics, Shanghai Jiaotong University, Shanghai 200240, China*

⁵*Center for Cold Atom Physics, Chinese Academy of Sciences, Wuhan 430071, China*

⁶*Department of Physics, Zhejiang Normal University, Jinhua 321004, China*



(Received 9 February 2018; revised manuscript received 10 March 2018; published 21 May 2018)

One of the outstanding challenges to information processing is the eloquent suppression of energy consumption in the execution of logic operations. The Landauer principle sets an energy constraint in deletion of a classical bit of information. Although some attempts have been made to experimentally approach the fundamental limit restricted by this principle, exploring the Landauer principle in a purely quantum mechanical fashion is still an open question. Employing a trapped ultracold ion, we experimentally demonstrate a quantum version of the Landauer principle, i.e., an equality associated with the energy cost of information erasure in conjunction with the entropy change of the associated quantized environment. Our experimental investigation substantiates an intimate link between information thermodynamics and quantum candidate systems for information processing.

DOI: [10.1103/PhysRevLett.120.210601](https://doi.org/10.1103/PhysRevLett.120.210601)

It was Landauer who expostulated, for the first time, a minimum amount of energy required to be consumed in deletion of a classical bit of information, known as the Landauer principle (LP) [1], implying an irreversibility of logical operations [2–4]. In terms of the LP, the erasure of information is fundamentally a dissipative process, which dissipates at least $k_B T \ln 2$ amount of heat, called the Landauer bound, from the system into the attached reservoir, where k_B and T represent the Boltzmann constant and reservoir temperature, respectively. On the other hand, if we try to understand violation of the second law of thermodynamics by Maxwell's demon (an intellectual creature envisioned by Maxwell), who converts thermal energy of the reservoir into useful work [5–7], the entropy cost should be considered regarding the demon's memory, which is also subject to the LP. So far, much of the discussion has been devoted to the validity and usefulness of the LP, including some skepticism and misunderstanding [8–12]. Since suppression of energy dissipation to a possible minimum is indispensable towards the continued development of digital computers [13,14], even for quantum information processing [15], further investigation of the LP is certainly of particular importance.

Some experimental attempts, subject to the conventional inequality of the LP, have been carried out to approach the Landauer bound using single-bit operations [16–20].

The latest experiment [20], for example, was implemented on a nanoscale digital magnetic memory bit, by which the intrinsic energy dissipation per single-bit operation was detected. In contrast, a very relevant investigation for extracting $k_B T \ln 2$ of heat in the creation of one bit of information [21], which is the complementary study of the LP in the context of a Szilárd engine [22]—a quantitative Maxwell demon—was accomplished recently on a single-electron transistor. However, all these aforementioned considerations surmise that the heat related to the removal or generation of a bit of information is regarding an open environment, which is a reservoir with much bigger size than the system. If we extend the treatments to a quantum domain, e.g., the information to be erased being encoded in a qubit and the environment being quantized, the model needs to be substantially reconsidered. The entropy of a quantum system is associated with the characteristic of the state, rather than the thermodynamic arrow of time. Additionally, in contrast to a large reservoir considered in the original version of the LP, the quantized reservoir is of a finite size and hence vacillating in interaction with the qubit system under quantum operations for information erasure. So, different from the previous assumption of final product states [23–25], quantum correlation appears between the system and the quantized reservoir during a quantum erasure process, which requires evaluation via mutual information and relative entropy. However, even with

elaborately designed systems under the consideration of a quantized reservoir, experimentally measuring mutual information and relative entropy is not easy to accomplish [26,27], which hampers efforts to exactly approach the Landauer bound.

Here we report an experimental investigation of a quantum mechanical LP by an experimental evaluation of system-reservoir correlation and entropy change during the erasure process. Our operations are based on a single ultracold $^{40}\text{Ca}^+$ ion confined in a linear Paul trap. Trapped ultracold ions, with the possibility of precise manipulation, have been considered as an ideal platform to explore the thermodynamics in a quantum domain with ultimate accuracy [28–31]. For our purpose, we consider the two internal levels of the ion as the qubit system and the vibrational degree of freedom of the ion as a finite-temperature reservoir. Removing information encoded initially in the qubit, we perceive the quantized LP by observing the phonon number variation in the quantized vibration of the ion. Owing to precise control of the 2 degrees of freedom in a coherent way, we demonstrate for the first time the possibility to approach the Landauer bound solely in a quantum mechanical fashion.

Our exploration mainly follows the idea termed as an improved LP [32], which describes the energy cost regarding information erasure employing an equality, rather than the conventional inequality. Assuming initial states of the system and reservoir to be a mixed state ρ_S and a thermal state ρ_R , correspondingly, our model starts from an uncorrelated state $\rho_{SR} = \rho_S \otimes \rho_R$ and evolves to a system-reservoir correlated state ρ'_{SR} resulting from an erasure process, where the system and reservoir states turn to $\rho'_S = \text{Tr}_R[\rho'_{SR}]$ and $\rho'_R = \text{Tr}_S[\rho'_{SR}]$, respectively. The improved LP gives a tight bound of the heat cost for an information erasure by an equality as

$$\Delta Q/k_B T = \Delta S + I(S':R') + D(\rho'_R||\rho_R), \quad (1)$$

where $\Delta Q = \text{Tr}[H_R \rho'_R] - \text{Tr}[H_R \rho_R]$ represents an average energy increase of the thermal reservoir with the Hamiltonian H_R . $\Delta S = S(\rho_S) - S(\rho'_S)$ denotes the entropy decrease of the system with the von Neumann entropy defined as $S(\hat{\rho}) = -\text{Tr}[\hat{\rho} \log \hat{\rho}]$. $I(S':R') = S(\rho'_S) + S(\rho'_R) - S(\rho'_{SR})$ indicates mutual information related to the system-reservoir correlations following the erasure process, and the relative entropy $D(\rho'_R||\rho_R) = \text{Tr}[\rho'_R \log \rho'_R] - \text{Tr}[\rho'_R \log \rho_R]$ corresponds to the free-energy increase of the reservoir during the erasure process. In comparison with the original form of the LP, i.e., $\Delta Q/k_B T \geq \Delta S$, Eq. (1) is obviously a quantum version of the LP linking quantum information with thermodynamics.

Our experiment is performed in a typical linear Paul trap with four parallel blade-like electrodes, and two end-cap electrodes aligned along the axial direction [33,34]. The trap is driven by the rf $\Omega_{\text{rf}}/2\pi = 17.6$ MHz with the power

of 5 W, giving the radial frequency $\omega_r/2\pi = 1.2$ MHz, and the trap axial frequency under the pseudopotential approximation is $\omega_z/2\pi = 1.01$ MHz with a voltage of 720 V applied to the end-cap electrodes. Under an external magnetic field of 6 G, the ground state $4^2S_{1/2}$ and the metastable state $3^2D_{5/2}$ split into two and six hyperfine energy levels, respectively, as shown in Fig. 1(a). We encode $|\downarrow\rangle$ in $|4^2S_{1/2}, m_J = +1/2\rangle$ and $|\uparrow\rangle$ in $|3^2D_{5/2}, m_J = +3/2\rangle$, where m_J is the magnetic quantum number. After Doppler cooling and optical pumping, a resolved sideband cooling is applied to cool the z -axis motional mode down to its vibrational ground state with the final average phonon number of 0.030(7), and the internal state of the ion is initialized to $|\downarrow\rangle$ with a probability of 98.9(2.3)%. The population of $|\downarrow\rangle$ could be detected by a 397-nm laser with detecting error $\leq 0.22(8)\%$ [35]. A narrow-linewidth 729-nm laser with wave vector \mathbf{k} radiates the ultracold ion with an incident angle of 22.5° with respect to the z axis, yielding the Lamb-Dicke parameter η of 0.09.

Figure 1(b) presents a schematic of our work. We start from preparing the qubit in an equal probability admixture of the states $|\downarrow\rangle$ and $|\uparrow\rangle$, indicating that the initial entropy of the system is superlative, but in contrast, our available information about the system is minimum; that is, we have no explicit information about the existing state of the ion. For a general case, we consider the reservoir to be initially in a thermal state with an associated energy defined as $E_0 = \text{Tr}[H_R \rho_R]$ for a given state ρ_R [see Fig. 1(b1)]. The erasure is performed by employing a red-sideband transition, governed by $H_R = \eta \hbar \Omega (a \sigma_+ e^{i\phi} + a^\dagger \sigma_- e^{-i\phi})/2$, where Ω and ϕ represent the Rabi frequency and laser phase, respectively, σ_+ (σ_-) indicates the raising (lowering) operator of the spin, and a (a^\dagger) denotes the annihilation (creation) operator of the quantized vibration of the ion in the z direction. The erasure process produced by $U_R = \exp(-iH_R t)$, with t as the pulse length, causes the population transfer from $|\uparrow\rangle$ to $|\downarrow\rangle$, corresponding to an entropy decrease of the system, which is accompanied by the reservoir energy increase up to E_m , as depicted in Fig. 1(b2). The erasure process ends with the system in the polarized state $|\downarrow\rangle$ and the reservoir changes to the final state ρ'_R , the energy of which is given as $E_f = \text{Tr}[H_R \rho'_R]$; see Fig. 1(b3). So we finally have a complete enlightenment about the state of the system accomplished by virtue of the heat increment $\Delta Q = E_f - E_0$. In this way, we may verify Eq. (1) by observing the variation of the population and phonon number in the qubit and the reservoir, respectively, where the entropy decrease of a system is $\Delta S = \ln 2$ in an ideal erasure for the system attached to a zero-temperature reservoir. The corresponding experimental procedure is drawn in Fig. 1(c).

In our actual implementation, superposition of the qubit states was achieved by a carrier-transition operator $U_C(\theta_c, 0) = \cos(\theta_c/2)I - i \sin(\theta_c/2)\sigma_x$, where σ_x is the Pauli operator. Since the dephasing rate (~ 0.5 kHz) of the




FIG. 1. A single trapped ion manipulated for exploring an improved LP. (a) Level scheme of the $^{40}\text{Ca}^+$ ion confined in a linear Paul trap under an external magnetic field. The qubit encoded in $|\uparrow\rangle$ and $|\downarrow\rangle$ represents the system, and the reservoir is denoted by the quantized z -axis vibration of the ion. The system-reservoir coupling is well controlled by red- or blue-sideband 729-nm laser radiation, and readout of the qubit is carried out by 397-nm spontaneous emission. (b) Illustrative schematic. (1) Starting point, where the qubit (i.e., the system) is in a maximally mixed state $\rho_S = (|\uparrow\rangle\langle\uparrow| + |\downarrow\rangle\langle\downarrow|)/2$ and the vibrational degree of freedom (i.e., the reservoir) is thermalized to energy E_0 . (2) Intermediate step, where the 729-nm laser drives a unitary evolution of the system and reservoir. The entropy of the system decreases, while the energy of the reservoir increases to E_m . (3) End of the erasure, where the system is populated in $|\downarrow\rangle$ and the reservoir ends at a higher level with energy E_f . (c) Flow diagram of the experimental operations, where (1) is to reach a mixed state of the system, (2) is related to a thermal state of the reservoir, (3) is to monitor the population of the system state, and (4) refers to measuring the average phonon number in the reservoir by observing the blue-sideband Rabi oscillation.

qubit is much larger than its decay rate (~ 1 Hz), waiting for a time longer than 2 ms would definitely lead to qubit dephasing but without affecting the population, which guarantees the removal of the off-diagonal terms of the density matrix of the aforementioned model while retaining the diagonal terms unaffected, yielding the initial state $\rho_S = \alpha|\downarrow\rangle\langle\downarrow| + \beta|\uparrow\rangle\langle\uparrow|$, with $\alpha = \cos^2(\theta_c/2)$ and $\beta = \sin^2(\theta_c/2)$. Meanwhile, the reservoir was thermalized due to switch-off of the cooling lasers during this waiting time. Consequently, our implementation commences with the reservoir at a finite temperature, where the average phonon number $\langle n \rangle_0$ is determined by waiting for a desired time duration. To remove the information from the system as much as we can, we have considered an optimal erasure under the first-order approximation where the largest contribution term affecting the erasure is eliminated during the red-detuned laser radiation for $t_{\text{op}} = \pi/\eta\Omega = 33(2) \mu\text{s}$ [34]. With this short-time implementation, decoherence effects, during the information removal, are negligible.

In our current study, all the four parts of Eq. (1) are explored experimentally as shown below. In our setup, the knowledge about the phonon number is obtained from qubit state measurement under the governance of the blue-detuning Hamiltonian $H_B = \eta\hbar\Omega(a\sigma_-e^{i\phi} + a^\dagger\sigma_+e^{-i\phi})/2$. In Figs. 2(a) and 2(b) we show, respectively, the

experimental observations of the blue-sideband Rabi oscillations versus the waiting time before and after the erasure, which is accomplished via the curve-fitting approach [36]. Consequently, based on the experimental data, the phonon number due to heat variation, as plotted in Fig. 2(c), could be calculated by Eq. (S14) in Ref. [34], and meanwhile the relative entropy of the reservoir is acquired. However, the populations in $|\downarrow\rangle$ and $|\uparrow\rangle$ at the end of the erasure are monitored from the 397-nm spontaneous-emission fluorescence spectrum [see Fig. 2(d)], which yields a variation of the system entropy. Moreover, the mutual information term is smaller with respect to other terms in Eq. (1), and a direct measurement of the mutual information is complicated due to its relevance to the system-reservoir entanglement. Therefore, we estimated the mutual information by an alternative way, as explained in Ref. [34], instead of direct experimental measurements.

With the data measured above, the Landauer bound and the improved LP are witnessed under the different reservoir temperatures in Fig. 2(e). Our observations clearly exhibit that the equality in the original form of the LP cannot be accessed quantum mechanically at the finite temperature. In contrary to the classical version of the LP in which the huge reservoir changes very little during the erasure process such that the mutual information and the relative entropy of the




FIG. 2. Experimental implementation of information erasure in a qubit system. (a) Blue-sideband Rabi oscillations before the erasure for different waiting time τ . (b) Blue-sideband Rabi oscillation at the end of erasure for different waiting time τ . (c) Average phonon number in the reservoir before the erasure (red dots) and after the erasure (black circles), obtained by calculating the data in (a) and (b). (d) Population in $|\downarrow\rangle$ (black circles) and $|\uparrow\rangle$ (red dots) at the end of erasure. The system is initialized with populations of 0.531(15) and 0.467(16) in $|\downarrow\rangle$ and $|\uparrow\rangle$, respectively. Dots or circles are experimental data and curves are analytical results. Each data point in (a) and (b) is measured by 10^2 repetition and in (d) by 10^4 repetition. (e) Test of the improved LP at different reservoir temperatures. The curves from top to bottom correspond to $\Delta Q/k_B T$ with $\Delta Q = Q_0(\langle n \rangle - \langle n \rangle_0)$ (black curve and crosses), $\Delta S + I(S':R') + D(\rho'_R||\rho_R)$ (red curve and circles), and ΔS (blue curve and dots), where the reservoir temperature is defined as $T = T_0/\ln[1 + 1/\langle n \rangle_0]$, with $\langle n \rangle_0$ the initial average phonon number. Inset of (e) demonstrates the result of $\Delta Q/k_B T - [\Delta S + I(S':R') + D(\rho'_R||\rho_R)]$. Experimental data in (e) are obtained from (a)–(d). The energy and temperature units are $Q_0 := \hbar\omega_z = 6.7 \times 10^{-28}$ J and $T_0 := Q_0/k_B = 48.5$ μ K, respectively. The curves in each panel are analytical results in our optimal erasure scheme using a π pulse of the red-sideband radiation [34]. The associated error bars reveal standard deviation.

reservoir can be ignored, both states of the system and reservoir in the quantum regime vary during the erasure process, which yields a quantum correlation between the system and the reservoir. Intrinsically, we can observe a large disparity between the heat consumption of the reservoir (black curves) and the entropy decrease of the system (blue curves), due to the difference from quantum correlation and the relative entropy of the reservoir. This difference turns out to be larger at lower temperature because in this case the relative entropy becomes more dominant in Eq. (1). In particular, in the limit of zero temperature, one may find the system's entropy decrease $\Delta S = \ln 2$, the mutual information $I(S':R') = 0$, and the relative entropy $D(\rho'_R||\rho_R) \rightarrow \infty$, which agrees with the result of $1/T \rightarrow \infty$ in the left-hand side of Eq. (1). In this scenario, the nearly perfect overlap between the red dashed and black solid curves implies that Eq. (1) provides a better way to understand the LP and the associated Landauer bound in the quantum regime.

It is interesting to note that the quantum LP is very sensitive to the initial condition of the model. Figure 3 presents the variation of the system entropy from negative to positive and then to negative again, arguing that entropy decrease, i.e., removal of information, occurs in the system for $\theta_c \in [0.54(3), 2.80(2)]$, and otherwise, the system entropy increases during the erasure, indicating creation of information. This indicates that our implementation could demonstrate both the generation and deletion of the system's information by simply tuning θ_c (i.e., varying the initial state of the system). Particularly, the generation of information in the region of $\theta_c < 0.54(3)$ corresponds to a fully quantized single-qubit Szilárd engine, the opposite process of the quantum LP. However, the observation in the region of $\theta_c > 2.80(2)$ is counterintuitive, where the generation of information is accompanied by the energy increase of the reservoir. This phenomenon is completely different from the aforementioned quantum LP and Szilárd engine, but reflects the significant role of the relative




FIG. 3. Experimental test of the improved LP under different initial states of the system. Curves from top to bottom correspond to $\Delta Q/k_B T$ (black curve and crosses), $\Delta S + I(S':R') + D(\rho'_R \parallel \rho_R)$ (red curve and circles), and ΔS (blue curve and dots), where the reservoir temperature is initially $18.2(8) \mu\text{K}$ corresponding to the average phonon number $\langle n \rangle_0 = 0.074(9)$. The horizontal dashed line indicates the positive-negative switch of the entropy decrease with boundary points at $\theta_c = 0.54(3)$ and $2.80(2)$. Inset demonstrates the result of $\Delta Q/k_B T - [\Delta S + I(S':R') + D(\rho'_R \parallel \rho_R)]$. Dots, circles, and crosses represent experimental data; curves are from analytical results. The system entropy is initially $S_1 = S\{\text{diag}[\cos^2(\theta_c/2), \sin^2(\theta_c/2)]\}$, with diag indicating a diagonal matrix and θ_c associated with the initial state of the system. The error bars indicate standard deviation.

entropy of the reservoir played in the fully quantized system-reservoir model [37].

In summary, we have demonstrated, in a more pertinent way, the first experimental investigation of the quantum mechanical LP using a single ultracold trapped ion. This is an imperative step towards better understanding of the fundamental physical limitations of irreversible logic operations at the quantum level. Our observation confirms that the LP still holds even at the quantum level after some modification by introducing quantum information quantities. Our experimental evidence might be helpful for efficient initialization of a future quantum computer involving an artificial quantum reservoir for fast elimination of the encoded information from large numbers of qubits, in which the information erasure yields more heat consumption than the classical counterpart due to system-reservoir correlations. For the same reason, quantum error correction in such a quantum computer would also cost higher heat (or energy). We believe that the experiment reported here will open an avenue towards further exploration of the Landauer bound in the quantum regime as well as possible applications.

This work was supported by National Key Research & Development Program of China under Grant

No. 2017YFA0304503, by National Natural Science Foundation of China under Grants No. 11734018, No. 11674360, No. 11404377, No. 91421111, No. 11571313, and No. 11574353, and by the Strategic Priority Research Program of the Chinese Academy of Sciences under Grant No. XDB21010100. T. P. X. was partially supported by Guangxi Natural Science Foundation of China (2015GXNSFAA139307), Guangxi Key Laboratory of Trusted Software (kx201505), Program for Innovative Research Team of Guilin University of Electronic Technology. K. R. thankfully acknowledges support from CAS-TWAS president's fellowship.

L. L. Y., T. P. X., and K. R. contributed equally to this work.

Note added.—Recently we became aware of a previous work [38] which presented an equality to be equivalent to Eq. (1) above, although no connection to Landauer principle was made explicitly there.

*zhoufei@wipm.ac.cn

†ywl@wipm.ac.cn

*mangfeng@wipm.ac.cn

- [1] R. Landauer, *IBM J. Res. Dev.* **5**, 183 (1961).
- [2] R. Landauer, *Nature (London)* **335**, 779 (1988).
- [3] S. Lloyd, *Nature (London)* **406**, 1047 (2000).
- [4] J. D. Meindl and J. A. Davis, *IEEE J. Solid-State Circuits* **35**, 1515 (2000).
- [5] O. Penrose, *Foundations of Statistical Mechanics: A Deductive Treatment* (Pergamon, New York, 1970).
- [6] C. H. Bennett, *Int. J. Theor. Phys.* **21**, 905 (1982).
- [7] K. Maruyama, F. Nori, and V. Vedral, *Rev. Mod. Phys.* **81**, 1 (2009).
- [8] O. J. E. Maroney, *Stud. Hist. Phil. M. P.* **36**, 355 (2005).
- [9] J. D. Norton, *Stud. Hist. Phil. M. P.* **42**, 184 (2011).
- [10] L. B. Kish and C. G. Granqvist, *Europhys. Lett.* **98**, 68001 (2012).
- [11] M. Hemmo and O. R. Shenker, *The Road to Maxwell's Demon: Conceptual Foundations of Statistical Mechanics* (Cambridge University Press, Cambridge, England, 2012).
- [12] J. D. Norton, *Entropy* **15**, 4432 (2013).
- [13] M. P. Frank, *Comput. Sci. Eng.* **4**, 16 (2002).
- [14] E. Pop, *Nano Res.* **3**, 147 (2010).
- [15] M. B. Plenio and V. Vitelli, *Contemp. Phys.* **42**, 25 (2001).
- [16] G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles, and D. J. Evans, *Phys. Rev. Lett.* **89**, 050601 (2002).
- [17] V. Blickle, T. Speck, L. Helden, U. Seifert, and C. Bechinger, *Phys. Rev. Lett.* **96**, 070603 (2006).
- [18] A. Béruit, A. Arakelyan, A. Petrosyan, S. Ciliberto, R. Dillenschneider, and E. Lutz, *Nature (London)* **483**, 187 (2012).
- [19] Y. Jun, M. Gavrilov, and J. Bechhoefer, *Phys. Rev. Lett.* **113**, 190601 (2014).
- [20] J. Hong, B. Lambson, S. Dhuey, and J. Bokor, *Sci. Adv.* **2**, e1501492 (2016).
- [21] J. V. Koski, V. F. Maisi, J. P. Pekola, and D. V. Averin, *Proc. Natl. Acad. Sci. U.S.A.* **111**, 13786 (2014).

- [22] L. Szilárd, in *Maxwell's Demon: Entropy, Information, Computation*, edited by H. S. Leff and A. E. Rex (Princeton University Press, Princeton, NJ, 1990).
- [23] B. Piechocinska, *Phys. Rev. A* **61**, 062314 (2000).
- [24] H. S. Leff and A. R. Rex, *Maxwell's Demon: Entropy, Classical and Quantum Information, Computing* (Institute of Physics Publishing, Bristol, 2003).
- [25] S. W. Kim, T. Sagawa, S. DeLiberato, and M. Ueda, *Phys. Rev. Lett.* **106**, 070401 (2011).
- [26] J. P. S. Peterson, R. S. Sarthour, A. M. Souza, I. S. Oliveira, J. Goold, K. Modi, D. O. Soares-Pinto, and L. C. Céleri, *Proc. R. Soc. A* **472**, 813 (2015).
- [27] B. Gaudenzi, E. Burzuri, S. Maegawa, H. S. J. van der Zant, and F. Luis, [arXiv:1703.04607v2](https://arxiv.org/abs/1703.04607v2).
- [28] S. An, J.-N. Zhang, M. Um, D. Lv, Y. Lu, J. Zhang, Z.-Q. Yin, H. T. Quan, and K. Kim, *Nat. Phys.* **11**, 193 (2015).
- [29] G. Huber, F. Schmidt-Kaler, S. Deffner, and E. Lutz, *Phys. Rev. Lett.* **101**, 070403 (2008).
- [30] O. Abah, J. Rossnagel, G. Jacob, S. Deffner, F. Schmidt-Kaler, K. Singer, and E. Lutz, *Phys. Rev. Lett.* **109**, 203006 (2012).
- [31] J. Rossnagel, O. Abah, F. Schmidt-Kaler, K. Singer, and E. Lutz, *Phys. Rev. Lett.* **112**, 030602 (2014).
- [32] D. Reeb and M. M. Wolf, *New J. Phys.* **16**, 103011 (2014).
- [33] F. Zhou, L. L. Yan, S. J. Gong, Z. H. Ma, J. Z. He, T. P. Xiong, L. Chen, W. L. Yang, M. Feng, and V. Vedral, *Sci. Adv.* **2**, e1600578 (2016).
- [34] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.120.210601> for details.
- [35] T. P. Xiong, L. L. Yan, F. Zhou, K. Rehan, D. F. Liang, L. Chen, W. L. Yang, Z. H. Ma, M. Feng, and V. Vedral, *Phys. Rev. Lett.* **120**, 010601 (2018).
- [36] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, *Phys. Rev. Lett.* **76**, 1796 (1996).
- [37] L. Rio, J. Åberg, R. Renner, O. Dahlsten, and V. Vedral, *Nature (London)* **474**, 61 (2011).
- [38] M. Esposito, K. Lindenberg, and C. Van den Broeck, *New J. Phys.* **12**, 013013 (2010).